

Gynecologic Oncology

An International Journal

EDITOR
DAVID M. GERSHENSON

Univ. of Minn.
Bio-Medical
Library

Associate Editors
Larry J. Copeland William J. Hoskins Karl C. Podratz

The Official Publication of
Society of Gynecologic Oncologists
Society of Gynecologic Oncologists of Canada
Felix Rutledge Society
Western Association of Gynecologic Oncologists
The Society of Memorial Gynecologic Oncologists
New England Association of Gynecologic Oncologists
Mid Atlantic Gynecologic Oncologic Society



ACADEMIC PRESS



Serum Levels of Soluble Intercellular Adhesion Molecule-1 (ICAM-1) and the Expression of ICAM-1 mRNA in Uterine Cervical Cancer

Kaei Nasu, M.D., Hisashi Narahara, M.D., Yasuko Etoh, M.D., Yasushi Kawano, M.D.,
Yoshiko Hirota, M.D., and Isao Miyakawa, M.D.

Department of Obstetrics and Gynecology, Oita Medical University, Hasama-machi, Oita 879-55, Japan

Received September 16, 1996

The purpose of this investigation was to evaluate the serum level of soluble intercellular adhesion molecule-1 (sICAM-1) in patients with squamous cell carcinoma of the cervix. Serum levels of sICAM-1 were measured by enzyme-linked immunosorbent assay in patients with cervical cancer (stage 0 through IV). Expression of mRNA in tumor tissues was evaluated by reverse transcriptase-polymerase chain reaction. Serum level of sICAM-1 in patients with advanced-stage (II-IV) or recurrent cervical cancer was found to be increased significantly. Expression of ICAM-1 mRNA was observed in most tumor tissues. Results suggest that sICAM-1 is shed from the cancerous tissue in patients with squamous cell carcinoma of the cervix. © 1997 Academic Press

Cell-matrix interactions are essential for the adhesion and migration of cells, wound healing, cell differentiation, and polarity [1, 2]. These interactions are mediated by the adhesion of cell-surface receptors, particularly of the integrins, to adhesive proteins [2, 3].

Intercellular adhesion molecule-1 (ICAM-1, CD54), a member of the immunoglobulin superfamily of adhesion receptors, has five extracellular immunoglobulin-like domains, a single transmembrane region, and a short cytoplasmic tail [4, 5]. ICAM-1 is expressed on various cell types, including fibroblasts, keratinocytes, vascular endothelial cells, macrophages, and glandular epithelial cells and stromal cells in the uterus [2, 6, 7]. ICAM-1 is a ligand for lymphocyte function-associated antigen-1 (LFA-1 or CD11a/CD18), which is present on lymphocytes, monocytes, and neutrophils [8].

ICAM-1 is an important early marker of immune activation and response [9-11]. The upregulated expression of ICAM-1 on cell surfaces is observed in a variety of diseases, including patients with rejection of liver allograft [9], autoimmune thyroiditis [10], Sjögren's syndrome [12], asthma [13], cutaneous T-cell lymphoma [14], melanoma [15], prostate cancer [16], ovarian cancer [17], and colonic cancer [18]. Coleman *et al.* [19] demonstrated the expression of

ICAM-1 protein in high-grade intraepithelial squamous neoplasia of the cervix by immunohistochemistry and suggested that the expression was related to human papillomavirus infection.

Seth *et al.* [20] used a chemiluminescence technique to detect soluble ICAM-1 (sICAM-1) in the serum obtained from healthy young volunteers. They found that ICAM-1 existed in a membranous and a soluble form that lacks the cytoplasmic tail and the transmembrane region. Although its functional characteristics have not been elucidated, Rothlein *et al.* [21] suggested that the circulating molecule reflects inflammation, tissue damage, and proteolysis. When increased sICAM-1 binds LFA-1, which is expressed on lymphocytes, the lymphocytes cannot bind other cells via adhesion molecules, resulting in the inhibition of lymphocyte function [22].

In the present study, we measured serum levels of sICAM-1 in patients with squamous cell carcinoma of the cervix, and investigated the expression of ICAM-1 in tumor tissues by reverse transcriptase-polymerase chain reaction (RT-PCR).

MATERIALS AND METHODS

ELISA for detection of sICAM-1. We obtained 6-ml serum samples from 34 Japanese women with squamous cell carcinoma of the cervix before the initial treatment (9 in stage 0, 7 in stage I, 3 in stage II, 5 in stage III, and 3 in stage IV), 7 patients with a recurrence of this cancer, and 16 healthy volunteers. Venous blood was allowed to clot at 4°C and was then centrifuged at 1000g for 15 min to remove the clot. Each serum sample was divided into 1-ml aliquots and frozen at -70°C until assayed.

Serum levels of sICAM-1 were measured by a sandwich enzyme-linked immunosorbent assay (ELISA) through use of an ICAM-1 test kit (T Cell Diagnostics, Cambridge, MA). The serum was allowed to thaw at room temperature and was then diluted 100-fold in a dilution buffer (T Cell Diag-

nostics). The assay was performed in duplicate as follows: standards or diluted samples (25 μ l) were placed in polystyrene microtiter wells that were precoated with a mouse anti-human ICAM-1 monoclonal antibody. A horseradish peroxidase-conjugated mouse anti-human ICAM-1 monoclonal antibody was immediately added to the wells, which were then incubated for 2 hr at room temperature. After the wells had been washed with phosphate-buffered saline (PBS, T Cell Diagnostics), a chromogen solution (T Cell Diagnostics) was added. The reaction was terminated by the addition of 2 *N* H₂SO₄ (Wako, Osaka, Japan) and absorbance was measured at 490 nm by use of a microplate reader (Model 3550, Bio-Rad Laboratories, Richmond, CA). All the reactions were performed at room temperature.

Data are the mean \pm SD. Difference were evaluated by Scheffe's *F* test using StatView 4.01 (Abacus Concepts, Berkeley, CA). A value of *P* < 0.05 was regarded as statistically significant.

RT-PCR for detection of ICAM-1 mRNA. Samples (100-mg) of cervical squamous cell carcinoma tissues were obtained from seven patients in an advanced stage (stage III) of the disease. As controls, we obtained specimens of normal cervical tissue from four women who had undergone hysterectomy for the evaluation of a benign disease. As positive controls, we used myelomonocytic cell line, HL-60, on which ICAM-1 expression had previously demonstrated [23, 24]. Each tissue sample was soaked in 1 ml of Isogen solution (Nippon Gene, Tokyo, Japan), homogenized by use of a Polytron (Type PT 10/35, Kinematica GmbH, Luzern, Switzerland), stored at room temperature for 5 min, and shaken vigorously for 15 sec after the addition of 0.2 ml of chloroform. The homogenates were centrifuged at 12,000g at 4°C for 15 min and then 0.5 ml of isopropanol was added to the aqueous phase. Each aliquot was stored at room temperature for 10 min and centrifuged at 12,000g at 4°C for 10 min. To the precipitate, 1 ml of 75% ethanol (Wako) was added. The aliquot was shaken vigorously and centrifuged at 12,000g at 4°C for 15 min. The precipitate was then dried briefly and dissolved in water.

ICAM-1 gene expression was analyzed by the RT-PCR method through use of an RNA PCR Kit with AMV RTase (Takara, Tokyo, Japan). RNA was reverse-transcribed into complementary DNA (cDNA). In brief, the reaction mixture (5 mM MgCl₂, 1 \times RNA PCR buffer [10 mM Tris-HCl, 50 mM KCl, pH 8.3], 1 mM dNTP mixture [dATP, dCTP, dGTP, dTTP] [Takara], RNase Inhibitor, 1 U/ μ l [Takara], reverse transcriptase, 0.25 U/ μ l [Takara], and 2.5 μ M random 9 mer [Takara]) was added to the RNA solution containing 1 μ g of total RNA, and the mixture was incubated at 30°C for 10 min. After incubation at 50°C for 25 min, the mixture was heated to 99°C for 5 min and then chilled on ice.

To perform the PCR assay, primer sets for ICAM-1 (upper primer: 5'-GTCCCCCTCAAAAGTCATCCTG-3' and lower primer: 5'-CCCTCGTCCTCTGCGGTCACAC-3') were designed according to the previously reported sequence [23, 24], synthesized by the phosphoramidite method on a DNA synthesizer (Model 8700, Biosearch, San Rafael, CA), and purified on Sephadex G50 columns (Pharmacia LKB Biotechnology, Uppsala, Sweden) and by high-performance liquid chromatography (HPLC). Forward and reverse primers were located within the open reading frame; the predicted size of the PCR product was 759 bp. The size of the RT-PCR product is apparently different from that of the PCR product of genomic DNA.

The cDNA transcribed from 1 μ g of total RNA was amplified on a thermal cycler (Model PJ2000, Perkin Elmer, Norwalk, CT) in a total volume of 80 μ l containing 4 mM MgCl₂, 1 \times RNA PCR buffer, 0.2 μ M of each primer, and 2.5 units of *Taq* DNA polymerase (Takara). PCR was carried out for 40 cycles with thermal cycler conditions for denaturing at 94°C for 3 min, annealing at 59°C for 2 min, and extension at 72°C for 2 min. The PCR products were separated by 1.2% agarose gel (Takara) electrophoresis and visualized by ethidium bromide (Takara) staining.

RESULTS

Serum levels of sICAM-1. Serum levels of sICAM-1 in healthy volunteers, in patients with cervical cancer in stages 0, I, II, III, and IV and in patients with recurrent tumor were 294 \pm 84, 274 \pm 39, 339 \pm 136, 554 \pm 131, 498 \pm 155, 451 \pm 108, and 503 \pm 111 ng/ml, respectively. Although the serum level of sICAM-1 in patients with early-stage cervical cancer (stage 0–I) did not increase, serum levels of ICAM-1 in patients with advanced-stage cervical cancer or recurrent cancer were significantly increased (control vs stage II, *P* < 0.05; control vs stage III, *P* < 0.05; control vs recurrence, *P* < 0.01; stage 0 vs stage II, *P* < 0.025; stage 0 vs stage III, *P* < 0.05; stage 0 vs recurrence, *P* < 0.025; Scheffe's *F* test) (Table 1).

Expression of ICAM-1 mRNA. As shown in Fig. 1, ICAM-1 mRNA expression had been observed in HL-60 cells as previously described [23, 24]. The expression of ICAM-1 mRNA was observed in six of seven tumor tissues. ICAM-1 mRNA expression in four samples of normal cervical tissue was below the detection level.

DISCUSSION

ICAM-1 expression has been observed on various cell types in normal, inflammatory, and tumor tissues, including fibroblasts, keratinocytes, vascular endothelial cells, lymphocytes, monocytes, glandular epithelial cells and stromal cells in the uterus, carcinomas of the gastrointestinal tract

TABLE 1

Serum Levels of sICAM-1 in Healthy Volunteers, in Patients with Cervical Cancer in Stages 0, I, II, III, and IV, and in Patients with Recurrent Tumor

Group	Number of patients	Serum sICAM-1 (ng/ml) ^a
Control	16	249 ± 84
Stage 0	9	274 ± 39
Stage I	7	339 ± 136
Stage II	3	554 ± 131*†
Stage III	5	498 ± 155*††
Stage IV	3	451 ± 108
Recurrence	7	503 ± 111**†

^a Data are the mean ± SD.

* $P < 0.05$, ** $P < 0.01$, vs control, Scheffe's F test.

† $P < 0.025$, †† $P < 0.05$, vs stage 0, Scheffe's F test.

[18, 25], ovarian cancer [17], and malignant melanoma [2, 6, 7, 26].

The involvement of ICAM-1 in the pathogenesis of malignant tumors has been examined, especially in malignant melanoma. Reports by Johnson *et al.* [27], Natali *et al.* [15], and Kageshita *et al.* [28] suggest that increased expression of ICAM-1 on melanoma cells may be positively correlated with a greater risk of metastasis. In contrast, evidence from Vanky *et al.* [29] suggests that expression of ICAM-1 on tumor cells may facilitate recognition by autologous lymphocytes and thus reduce metastasis. Harning *et al.* [26] observed increased serum levels of sICAM-1 in patients with malignant melanoma and found a correlation between increasing serum level of sICAM-1 and prognosis. ICAM-1 is also expressed on keratinocytes and blood vessels in benign inflammatory disorders, such as dermatitis and allergic eczema, and this expression appears to be correlated with the extent of disease [14].

Tsujiaki *et al.* [25] used immunochemical methods to detect cell surface and sICAM-1 in cell lines obtained from colonic carcinoma, gastric carcinoma, pancreatic carcinoma, hepatocellular carcinoma, and lung adenocarcinoma. They also reported that serum levels of sICAM-1 were elevated in patients with gastric cancer, colonic cancer, gall bladder cancer, pancreatic cancer, and esophageal cancer, especially in patients with liver metastasis. They suggested that sICAM-1 may block the attachment of cytotoxic T cells and/or NK cells to cancer cells, since LFA-1 could be blocked with sICAM-1, and that the higher levels of serum sICAM-1 in malignant diseases may reflect the host's immune response to malignant cells and surrounding cells. Harning *et al.* [26] suggested that the elevated serum levels of ICAM-1 observed in patients with malignant melanoma are indicative of an enhanced host cell-mediated immune response to a primary tumor or may represent ICAM-1 shed *in situ* by

melanoma cells. Burrows *et al.* [30] maintained that ICAM-1 expression in most carcinomas is a passive event resulting from induction by infiltrating leukocytes and is unrelated to the tumor cells themselves. Yamamoto *et al.* [31] reported that peripheral blood mononuclear cells from cancer patients revealed an increase of ICAM-1 expression.

Coleman and Stanley [32] reported that the expressions of ICAM-1, vascular cellular adhesion molecule-1 (VCAM-1), and E-selectin were significantly up-regulated in the high-grade intraepithelial squamous neoplasia of the cervix using immunohistochemical method. ICAM-1 expression was observed in vessels, keratinocytes, and mononuclear cells in these lesions. They suggested that the enhanced expression of these adhesion molecules may be functionally important in enabling the local recruitment of immunocompetent cells.

Rothlein *et al.* [21] suggested that sICAM-1 serves as a marker of inflammation, tissue damage, and proteolysis. The source and function of sICAM-1 in humans remain unclear. Cardiac transplant rejection is associated with an increase in the level of sICAM-1, a finding that suggests an increase in sICAM-1 may indicate activation of cells that participate in allograft rejection [33]. sICAM-1 is also detected in cell culture supernatants of normal lymphocytes and established cancer cell lines [25, 34].

ICAM-1 expression in tissues is induced by various inflammatory cytokines, such as IL-1, TNF- α , and IFN- γ [6, 35]. These cytokines, which are produced by various cells, including inflammatory cells and tumor cells, sometimes upregulate cellular binding between tumor cells and endothelial cells through the induction of adhesion molecules [30] and increase the incidence of metastasis [36, 37].

In the present study, we showed that the serum level of sICAM-1 is significantly elevated in patients with advanced cervical squamous cell carcinoma and that ICAM-1 mRNA is expressed in tumor tissues. Therefore, an increase in the serum levels of sICAM-1 may reflect the enhanced expression of the molecule in the cancerous tissues. This increase in production is detected using ELISA perhaps due to the fact that the tumor reaches a certain size or spreads beyond the uterus (i.e., higher than or equal to stage II). Although sICAM-1 level of patients with stage IV disease did not have a significant difference compared to the patients with earlier stages, we think that the number of the patients with stage IV disease is too small to make a statistical analysis. Inflammatory cells may infiltrate the tumor tissue partly by



FIG. 1. Electrophoresis of RT-PCR products. Representative results are shown. Lane 1, normal cervical tissue; lanes 2–5, squamous cell carcinoma of the cervix (stage III); lane 6, HL-60 cells.

interaction of their membranous LFA-1 and ICAM-1 on endothelial cells. We hypothesize that sICAM-1 is produced mainly by carcinoma cells and that sICAM-1 suppresses the infiltration of inflammatory cells by a competitive inhibition of the membranous ICAM-1/LFA-1 interaction. Our results suggest that ICAM-1 is present in squamous cell carcinoma of the cervix.

ACKNOWLEDGMENT

This work was supported in part by Ministry of Education, Science, and Culture of Japan Grant-in-Aid 08771342 for Scientific Research.

REFERENCES

1. Ruoslahti E, Pierschbacher MD: New perspectives in cell adhesion: RGD and integrins. *Science* 238:491-497, 1987
2. Larson RS, Springer TA: Structure and function of leukocyte integrins. *Immunol Rev* 114:181-217, 1990
3. Nasu K, Ishida T, Setoguchi M, Higuchi Y, Akizuki S, Yamamoto S: Expression of wild-type and mutated rabbit osteopontin in *Escherichia coli*, and their effects on adhesion and migration of P388D1 cells. *Biochem J* 307:257-265, 1995
4. Springer TA: Adhesion receptors of the immune system. *Nature* 346:425-434, 1990
5. Staunton DE, Dustin ML, Erickson HP, Springer TA: The arrangement of the immunoglobulin-like domains of ICAM-1 and the binding sites for LFA-1 and rhinovirus. *Cell* 61:243-254, 1990
6. Dustin ML, Rothlein R, Bhan AK, Dinarello CA, Springer TA: Induction by IL-1 and interferon- γ : Tissue distribution, biochemistry and function of a natural adherence molecule (ICAM-1). *J Immunol* 137:245-254, 1986
7. Tabibzadeh SS, Poubouridis D: Expression of leukocyte adhesion molecules in human endometrium. *Am J Clin Pathol* 93:183-189, 1990
8. Marlin SD, Springer TA: Purified intercellular adhesion molecule-1 (ICAM-1) is a ligand for lymphocyte function-associated antigen 1 (LFA-1). *Cell* 51:813-819, 1987
9. Adams DH, Hutcheson SC, Shaw J, Rothlein R: Intercellular adhesion molecule-1 on allografts during rejection. *Lancet* 8672:1122-1125, 1989
10. Westman AP, Cohen S, Makgoba MW, Borysiewicz LK: Expression of an intercellular adhesion molecule-1, ICAM-1, by human thyroid cells. *J Endocrinol* 122:185-191, 1989
11. Vogetseder W, Feichinger H, Schulz TF, Schwaebler W, Tabaczewski P, Mitroic M, Böck G, Marth C, Dapunt O, Mikuz G, Dierich MP: Expression of 7F7-antigen, a human adhesion molecule identical to intercellular adhesion molecule-1, in human carcinomas and their stromal fibroblasts. *Int J Cancer* 43:768-773, 1989
12. Saito I, Terauchi K, Shimizu M, Nishimura S, Yoshino K, Takeuchi T, Tsubota K, Miyasaka N: Expression of cell adhesion molecules in the salivary and lacrimal glands of Sjögren's syndrome. *J Clin Lab Anal* 7:180-187, 1993
13. Wegner CD, Gundel RH, Reilly P, Haynes N, Letts LG, Rothlein R: Intercellular adhesion molecule-1 (ICAM-1) in the pathogenesis of asthma. *Science* 247:456-459, 1990
14. Vejlsøgaard GL, Ralfkiaer E, Avnstorp C, Czajkowski M, Marlin SD, Rothlein R: Kinetics and characterization of intercellular adhesion molecule-1 (ICAM-1) expression on keratinocytes in various inflammatory skin lesions and malignant cutaneous lymphomas. *J Am Acad Dermatol* 20:782-790, 1989
15. Natali P, Nicotra MR, Cavallere R, Bigotti A, Romano G, Temponi M, Ferrone S: Differential expression of intercellular adhesion molecule 1 in primary and metastatic melanoma lesions. *Cancer Res* 50:1271-1278, 1990
16. Wolff JM, Stephenson RN, Chisholm GD, Habib FK: Levels of circulating intercellular adhesion molecule-1 in patients with metastatic cancer of the prostate and the benign prostatic hyperplasia. *Eur J Cancer* 31A:339-341, 1995
17. Gardner MJ, Jones LMH, Causall JB, Turner GA: Expression of cell adhesion molecules on ovarian tumour cell lines and mesothelial cells, in relation to ovarian cancer metastasis. *Cancer Lett* 91:229-234, 1995
18. Dippold W, Wirtig B, Schwaebler W, Mayer W, Meyer zum Büschenfelde K-H: Expression of intercellular adhesion molecule 1 (ICAM-1, CD54) in colonic epithelial cells. *Gut* 34:1593-1597, 1993
19. Coleman N, Greenfield IM, Hare J, Kruger-Gray H, Chain BM, Stanley MA: Characterization and functional analysis of the expression of intercellular adhesion molecule-1 in human papillomavirus-related disease of cervical keratinocytes. *Am J Pathol* 143:355-367, 1993
20. Seth R, Raymond FD, Makgoba MW: Circulating ICAM-1 isoforms: Diagnostic prospectus for inflammatory and immune disorders. *Lancet* 338:83-84, 1991
21. Rothlein R, Mainolfi EA, Czajkowski M, Marlin SD: A form of circulating ICAM-1 in human serum. *J Immunol* 147:3788-3793, 1991
22. Becker JC, Dummer R, Harman AA, Burg G, Schmidt RE: Shedding of ICAM-1 from human melanoma cell lines induced by INF- γ and tumor necrosis factor- α : Functional consequences on cell-mediated cytotoxicity. *J Immunol* 147:4398-4401, 1991
23. Simmons D, Makgoba MW, Seed B: ICAM-1, an adhesion ligand of LFA-1, is homologous to the neural cell adhesion molecule NCAM. *Nature* 331:624-627, 1988
24. Staunton DE, Marlin SD, Surdowa C, Dustin ML, Springer TA: Primary structure of ICAM-1 demonstrates interaction between members of the immunoglobulin and integrin supergene families. *Cell* 52:925-933, 1988
25. Tsujisaki M, Imai K, Hirata H, Hanzawa Y, Masuya J, Nakano T, Sugiyama T, Matsui M, Hinoda Y, Yachi A: Detection of circulating intercellular adhesion molecule-1 antigen in malignant diseases. *Clin Exp Immunol* 85:3-8, 1991
26. Harning R, Mainolfi E, Bystryn J-C, Henn M, Merluzzi VJ, Rothlein R: Serum levels of circulating intercellular adhesion molecule 1 in human malignant melanoma. *Cancer Res* 51:5003-5005, 1991
27. Johnson JP, Stadel BG, Holzmann E, Riethmüller G: *De novo* expression of intercellular-adhesion molecule 1 in melanoma correlates with increased risk of metastasis. *Proc Natl Acad Sci USA* 86:641-644, 1989
28. Kageshita T, Nakamura T, Yamada M, Kuriya N, Arao T, Ferrone S: Differential expression of melanoma-associated antigens in acral lentiginous melanoma and in nodular melanoma lesions. *Cancer Res* 51:1726-1732, 1991
29. Vanky F, Wang P, Patarroyo M, Klein E: Expression of adhesion molecule ICAM-1 and major histocompatibility complex class I antigens on human cells is required for their interaction with autologous lymphomas in vitro. *Cancer Immunol Immunother* 31:19-27, 1990
30. Burrows FJ, Huskard DO, Hart IR, Marshall JF, Selkirk S, Poole S: Influence of tumor-derived interleukin 1 on melanoma-endothelial cell interactions in vitro. *Cancer Res* 51:4768-4775, 1991

31. Yamamoto T, Yoneda K, Ueta E, Osaki T: Serum cytokines, interleukin-2 receptor, and soluble intercellular adhesion molecule-1 in oral disorders. *Oral Surg Oral Med Oral Pathol* 78:727-735, 1994
32. Coleman N, Stanley MA: Characterization and functional analysis of the expression of vascular adhesion molecules in human papillomavirus-related disease of the cervix. *Cancer* 74:884-892, 1994
33. Ballantyne C, Mainolfi EA, Young JB, Windsor NT, Cocanougher B, Lawrence EC, Anderson DC, Rothlein R: Prognostic value of increased levels of circulating intercellular adhesion molecule-1 after heart transplant. *Clin Res* 39:285a, 1991
34. Faull RJ, Russ GR: Adhesion of lymphocytes to stimulated vascular endothelial cells occurs via ICAM-1-dependent and ICAM-1-independent pathways. *Transplant Proc* 22:2099-2100, 1990
35. Rothlein R, Czajkowski M, O'Neill MM, Marlin SD, Mainolfi E, Merluzzi VJ: Induction of intercellular adhesion molecule-1 on primary and continuous cell lines by pro-inflammatory cytokines. *J Immunol* 141:1665-1669, 1988
36. Giavuzzi R, Garofalo A, Bani MR, Abbate M, Ghezzi P, Boraschi D, Mantovani A, Dejana E: Interleukin 1-induced augmentation of experimental metastases from a human melanoma. *Cancer Res* 50:4771-4775, 1990
37. Kelly SA, Gschmeissner S, East N, Balkwill R: Enhancement of metastatic potential by γ -interferon. *Cancer Res* 51:4020-4027, 1991